

BIG GAIN®

ELITE 37:0 R400

XP #1310RBI

SUPPLEMENT FOR CATTLE FED IN CONFINEMENT FOR SLAUGHTER

MEDICATED

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT		400 gm/ton (200 mg/lb)
Monsensin		
GUARANTEED ANALYSIS		
Crude Protein, min	37.00%
Crude Fat, min	1.50%
Crude Fiber, max	5.00%
Calcium, min	3.50%
Calcium, max	4.50%
Phosphorus, min	0.60%
Magnesium, min	0.50%
Potassium, min	1.50%
Salt, min	4.50%
Sulfur, min	5.50%
Salt, max	0.60%
Selenium, min	5.00 ppm
Vitamin A, min	40,000 IU/lb.
Vitamin D, min	8,000 IU/lb.
Vitamin E, min	50 IU/lb.

INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Animal Protein Products, Calcium Carbonate, Salt, Magnesium Oxide, Calcium Sulfate, Potrome Animal Fat, (preserved with BHA), Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Manganese Oxide, Zinc Oxide, Mineral Oil, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Vegetable Oil.

FEEDING DIRECTIONS

Thoroughly mix at a rate of 1.0 to 1.5 pounds per head per day (6.0 to 6.50% of intake) to supply 0.14 to 0.42 mg/lb of bodyweight per day up to 480 mg/head/day of monensin. Or, thoroughly mix 125 (25 g/ton monensin) to 150 (30 g/ton monensin) pounds of Elite 37:0 with Monensin (type B) along with the remainder of ingredients to obtain one ton (2,000 lbs.) of a type C feed. Consult your Big Gain representative for complete feeding directions.

Caution: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended in the feeding directions. Do not feed to lactating goats. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

Warning: A withdrawal time has not been established for pre-nursing calves. Do not use in calves to be processed for veal.

Warning: Contains Copper sources. Do not expose to sheep or related species.

D-1

2/20/23

Manufactured by
BIG GAIN INC.
Mankato, MN 56001
NET WT 50 LB (22.68 KG) OR BULK INVOICE



This feed was produced in a facility certified in the American Feed Industry Association's Safe Feed/Safe Food Certification Program.

BIG GAIN®

ELITE 37:0 R400

XP #1310RBI

SUPPLEMENT FOR CATTLE FED IN CONFINEMENT FOR SLAUGHTER

MEDICATED

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT		400 gm/ton (200 mg/lb)
Monsensin		
GUARANTEED ANALYSIS		
Crude Protein, min	37.00%
Crude Fat, min	1.50%
Crude Fiber, max	5.00%
Calcium, min	3.50%
Calcium, max	4.50%
Phosphorus, min	0.60%
Magnesium, min	0.50%
Potassium, min	1.50%
Salt, min	4.50%
Sulfur, min	5.50%
Salt, max	0.60%
Selenium, min	5.00 ppm
Vitamin A, min	40,000 IU/lb.
Vitamin D, min	8,000 IU/lb.
Vitamin E, min	50 IU/lb.

INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Animal Protein Products, Calcium Carbonate, Salt, Magnesium Oxide, Calcium Sulfate, Potrome Animal Fat (preserved with BHA), Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Manganese Oxide, Zinc Oxide, Mineral Oil, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Vegetable Oil.

FEEDING DIRECTIONS

Thoroughly mix at a rate of 1.0 to 1.5 pounds per head per day (6.0 to 6.50% of intake) to supply 0.14 to 0.42 mg/lb of bodyweight per day up to 480 mg/head/day of monensin. Or, thoroughly mix 125 (25 g/ton monensin) to 150 (30 g/ton monensin) pounds of Elite 37:0 with Monensin (type B) along with the remainder of ingredients to obtain one ton (2,000 lbs.) of a type C feed. Consult your Big Gain representative for complete feeding directions.

Caution: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended in the feeding directions. Do not feed to lactating goats. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

Warning: A withdrawal time has not been established for pre-nursing calves. Do not use in calves to be processed for veal.

Warning: Contains Copper sources. Do not expose to sheep or related species.

D-1

2/20/23

Manufactured by
BIG GAIN INC.
Mankato, MN 56001
NET WT 50 LB (22.68 KG) OR BULK INVOICE



This feed was produced in a facility certified in the American Feed Industry Association's Safe Feed/Safe Food Certification Program.

BIG GAIN®

ELITE 37:0 R400

XP #1310RBI

SUPPLEMENT FOR CATTLE FED IN CONFINEMENT FOR SLAUGHTER

MEDICATED

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT		400 gm/ton (200 mg/lb)
Monsensin		
GUARANTEED ANALYSIS		
Crude Protein, min	37.00%
Crude Fat, min	1.50%
Crude Fiber, max	5.00%
Calcium, min	3.50%
Calcium, max	4.50%
Phosphorus, min	0.60%
Magnesium, min	0.50%
Potassium, min	1.50%
Salt, min	4.50%
Sulfur, min	5.50%
Salt, max	0.60%
Selenium, min	5.00 ppm
Vitamin A, min	40,000 IU/lb.
Vitamin D, min	8,000 IU/lb.
Vitamin E, min	50 IU/lb.

INGREDIENTS

Plant Protein Products, Processed Grain By-Products, Animal Protein Products, Calcium Carbonate, Salt, Magnesium Oxide, Calcium Sulfate, Potrome Animal Fat, (preserved with BHA), Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Manganese Oxide, Zinc Oxide, Mineral Oil, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Vegetable Oil.

FEEDING DIRECTIONS

Thoroughly mix at a rate of 1.0 to 1.5 pounds per head per day (6.0 to 6.50% of intake) to supply 0.14 to 0.42 mg/lb of bodyweight per day up to 480 mg/head/day of monensin. Or, thoroughly mix 125 (25 g/ton monensin) to 150 (30 g/ton monensin) pounds of Elite 37:0 with Monensin (type B) along with the remainder of ingredients to obtain one ton (2,000 lbs.) of a type C feed. Consult your Big Gain representative for complete feeding directions.

Caution: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended in the feeding directions. Do not feed to lactating goats. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

Warning: A withdrawal time has not been established for pre-nursing calves. Do not use in calves to be processed for veal.

Warning: Contains Copper sources. Do not expose to sheep or related species.

D-1

2/20/23

Manufactured by
BIG GAIN INC.
Mankato, MN 56001
NET WT 50 LB (22.68 KG) OR BULK INVOICE



This feed was produced in a facility certified in the American Feed Industry Association's Safe Feed/Safe Food Certification Program.