

BIG GAIN®

GAIN MASTER 55:35

P #1607

SUPPLEMENT FOR CATTLE FED IN CONFINEMENT FOR SLAUGHTER

MEDICATED

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT	
Monensin	450 gm/ton (225 mg/lb)
GUARANTEED ANALYSIS	
Crude Protein, min (includes not more than 35% crude protein equivalent from non-protein nitrogen)	55.00%
Crude Fat, min	1.00%
Crude Fiber, max	5.00%
Calcium, min	11.00%
Calcium, max	12.50%
Phosphorus, min	0.60%
Magnesium, min	1.00%
Potassium, min	2.85%
Salt, min	5.75%
Salt, max	6.75%
Sulfur, min	0.95%
Vitamin A, min	50,000 IU/lb.
Vitamin D, min	10,000 IU/lb.
Vitamin E, min	50 IU/lb.

INGREDIENTS

Plant Protein Products, Calcium Carbonate, Urea, Salt, Processed Grain By-Products, Potassium Chloride, Calcium Sulfate, Animal Protein Products, Magnesium Oxide, Monocalcium Phosphate, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Manganese Oxide, Zinc Oxide, Mineral Oil, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Porcine Animal Fat (preserved with BHA), Urea Formaldehyde Condensation Polymer, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement.

FEEDING DIRECTIONS

Thoroughly mix at a rate of 0.85 to 1.40 lb/head/day depending on the body weight of the cattle with roughages and feed continuously to supply 0.14 to 0.42 mg/lb of bodyweight per day up to 480 mg/head/day of monensin. Consult your Big Gain representative for complete feeding directions.

Caution: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding unutilized or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended in the feeding directions. Do not feed to lactating goats. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

Warning: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

D-2

10/25/23

Manufactured by
BIG GAIN INC.
Mankato, MN 56001
NET WT 50 LB (22.68 KG) OR BULK INVOICE



This feed was produced in a facility certified in the American Feed Industry Association's Safe Feed/Safe Food Certification Program.

BIG GAIN®

GAIN MASTER 55:35

P #1607

SUPPLEMENT FOR CATTLE FED IN CONFINEMENT FOR SLAUGHTER

MEDICATED

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT	
Monensin	450 gm/ton (225 mg/lb)
GUARANTEED ANALYSIS	
Crude Protein, min (includes not more than 35% crude protein equivalent from non-protein nitrogen)	55.00%
Crude Fat, min	1.00%
Crude Fiber, max	5.00%
Calcium, min	11.00%
Calcium, max	12.50%
Phosphorus, min	0.60%
Magnesium, min	1.00%
Potassium, min	2.85%
Salt, min	5.75%
Salt, max	6.75%
Sulfur, min	0.95%
Vitamin A, min	50,000 IU/lb.
Vitamin D, min	10,000 IU/lb.
Vitamin E, min	50 IU/lb.

INGREDIENTS

Plant Protein Products, Calcium Carbonate, Urea, Salt, Processed Grain By-Products, Potassium Chloride, Calcium Sulfate, Animal Protein Products, Magnesium Oxide, Monocalcium Phosphate, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Manganese Oxide, Zinc Oxide, Mineral Oil, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Porcine Animal Fat (preserved with BHA), Urea Formaldehyde Condensation Polymer, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement.

FEEDING DIRECTIONS

Thoroughly mix at a rate of 0.85 to 1.40 lb/head/day depending on the body weight of the cattle with roughages and feed continuously to supply 0.14 to 0.42 mg/lb of bodyweight per day up to 480 mg/head/day of monensin. Consult your Big Gain representative for complete feeding directions.

Caution: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding unutilized or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended in the feeding directions. Do not feed to lactating goats. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

Warning: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

D-2

10/25/23

Manufactured by
BIG GAIN INC.
Mankato, MN 56001
NET WT 50 LB (22.68 KG) OR BULK INVOICE



This feed was produced in a facility certified in the American Feed Industry Association's Safe Feed/Safe Food Certification Program.

BIG GAIN®

GAIN MASTER 55:35

P #1607

SUPPLEMENT FOR CATTLE FED IN CONFINEMENT FOR SLAUGHTER

MEDICATED

For the prevention and control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENT	
Monensin	450 gm/ton (225 mg/lb)
GUARANTEED ANALYSIS	
Crude Protein, min (includes not more than 35% crude protein equivalent from non-protein nitrogen)	55.00%
Crude Fat, min	1.00%
Crude Fiber, max	5.00%
Calcium, min	11.00%
Calcium, max	12.50%
Phosphorus, min	0.60%
Magnesium, min	1.00%
Potassium, min	2.85%
Salt, min	5.75%
Salt, max	6.75%
Sulfur, min	0.95%
Vitamin A, min	50,000 IU/lb.
Vitamin D, min	10,000 IU/lb.
Vitamin E, min	50 IU/lb.

INGREDIENTS

Plant Protein Products, Calcium Carbonate, Urea, Salt, Processed Grain By-Products, Potassium Chloride, Calcium Sulfate, Animal Protein Products, Magnesium Oxide, Monocalcium Phosphate, Manganese Sulfate, Zinc Sulfate, Copper Sulfate, Manganese Oxide, Zinc Oxide, Mineral Oil, Calcium Iodate, Cobalt Carbonate, Sodium Selenite, Porcine Animal Fat (preserved with BHA), Urea Formaldehyde Condensation Polymer, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement.

FEEDING DIRECTIONS

Thoroughly mix at a rate of 0.85 to 1.40 lb/head/day depending on the body weight of the cattle with roughages and feed continuously to supply 0.14 to 0.42 mg/lb of bodyweight per day up to 480 mg/head/day of monensin. Consult your Big Gain representative for complete feeding directions.

Caution: Do not allow horses or other equines access to feeds containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding unutilized or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended in the feeding directions. Do not feed to lactating goats. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

Warning: A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

D-2

10/25/23

Manufactured by
BIG GAIN INC.
Mankato, MN 56001
NET WT 50 LB (22.68 KG) OR BULK INVOICE



This feed was produced in a facility certified in the American Feed Industry Association's Safe Feed/Safe Food Certification Program.