



Lime Ridge Ag Supply Inc
LRAS 16% CALF #0220

MEDICATED

For 200 lb to 500 lb calves.

For the prevention and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii* in calves (excluding veal calves).

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium)..... 44.0 g/ton

Guaranteed Analysis

Crude Protein ,(Min).....	15.9	%
Crude Fat ,(Min).....	3.8	%
Crude Fiber ,(Max).....	6.0	%
ADF ,(Max).....	6.9	%
Calcium ,(Min).....	0.7	%
Calcium ,(Max).....	0.9	%
Phosphorus ,(Min).....	0.3	%
Selenium ,(Min).....	1	ppm
Vitamin A ,(Min).....	12,624.0	IU/lb

List of Ingredients

Grain products, plant protein products, soybean oil, vitamin E supplement, propionic acid, potassium sulfate, magnesium sulfate, bentonite, sodium selenite, salt, molasses products, vitamin A supplement, vitamin D3 supplement, zinc sulfate, manganous sulfate, copper sulfate, cobalt carbonate, calcium iodate, manganous oxide, zinc oxide, processed grain by-products, propionic acid (a preservative), cane molasses, vegetable oil refinery lipid (feed grade), lecithin, phosphoric acid, sulfuric acid, natural and artificial flavoring, propionic acid (preservative) and sodium benzoate (preservative), processed grain-by products, calcium carbonate, monocalcium phosphate.

Feeding Directions

Feed 5 to 6 pounds per head per day.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not feed undiluted. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not exceed the levels of Monensin recommended by the feeding directions, as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: For calves, excluding veal calves, feed at a rate of 0.14 to 1.0 milligrams Monensin per pound of body weight per day, depending upon the severity of challenge, up to maximum of 200 milligrams per head per day.

Manufactured by:

Lime Ridge Ag Supply, Inc
115 Minor St
PO Box 115
Lime Ridge, WI 53942
608-986-2626

Texturized

Net weight 50 lbs (22.67 kg)

MFG DATE: 04/08/2025

MEDICATED