



# Lime Ridge Ag Supply Inc

## LRAS 18% CALF #0226

### MEDICATED

For dairy replacement calves, 1 week to 3 months of age.

Calves excluding veal calves: For prevention and control of coccidiosis due to *Eimeria bovis* and *E. zuernii*. Prevents the development of house flies, stable flies, face flies, and horn flies in the manure of treated cattle.

ACTIVE DRUG INGREDIENT	
Monensin (as Monensin Sodium) .....	35 g/ton
ACTIVE INGREDIENT	
Diflubenzuron .....	4.56 mg/lb

### Guaranteed Analysis

Crude Protein ,(Min).....	18.0	%
Crude Fat ,(Min).....	4.1	%
Crude Fiber ,(Max).....	5.0	%
ADF ,(Max).....	6.3	%
Calcium ,(Min).....	0.8	%
Calcium ,(Max).....	1.1	%
Phosphorus ,(Min).....	0.4	%
Selenium ,(Min).....	0.9	ppm
Vitamin A ,(Min).....	8,865.5	IU/lb

### List of Ingredients

Grain products, Oat Hulls, Roughage products, soybean oil, diflubenzuron, Sodium Sulfate, Sodium Sulfite, Potassium Sulfate, Sodium Bentonite, Flavoring, potassium sulfate, magnesium sulfate, bentonite, vitamin E supplement, sodium selenite, salt, molasses products, vitamin A supplement, vitamin D3 supplement, zinc sulfate, manganous sulfate, copper sulfate, cobalt carbonate, calcium iodate, manganous oxide, zinc oxide, processed grain by-products, propionic acid (a preservative), cane molasses, vegetable oil refinery lipid (feed grade), lecithin, phosphoric acid, sulfuric acid, natural and artificial flavoring, propionic acid (preservative) and sodium benzoate (preservative), plant protein products, processed grain-by products, calcium carbonate, monocalcium phosphate.

### Feeding Directions

**Limitations: Feed at a rate of 0.14 to 1.0 milligrams monensin per pound of body weight per day, depending upon the severity of challenge, up to a maximum of 200 milligrams per head per day.**  
**Feed up to 6 lbs per head per day.**

**CAUTION:** Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

Do not allow horses or other equines access to feeds containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

**WARNING:** A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal.

**Manufactured by:**  
**Lime Ridge Ag Supply, Inc**  
**115 Minor St**  
**PO Box 115**  
**Lime Ridge, WI 53942**  
**608-986-2626**

Texturized  
**Net weight 50 lbs (22.67 kg)**

**MFG DATE: 07/22/2025**

**MEDICATED**