



Lime Ridge Ag Supply Inc
LRAS STEER GROWER #0842

MEDICATED

For beef cattle and/or replacements.

Growing cattle on pasture or in dry lot (stocker and feeder cattle and dairy and beef replacement heifers): For increased rate of weight gain. For prevention and control of coccidiosis caused by Eimeria bovis and Eimeria zuernii.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin sodium) 32.5 g/ton

Guaranteed Analysis

Crude Protein* ,(Min).....	11.9	%
Crude Fat ,(Min).....	3.5	%
Crude Fiber ,(Max).....	5.0	%
Calcium ,(Min).....	0.7	%
Calcium ,(Max).....	1.1	%
Phosphorus ,(Min).....	0.3	%
Potassium ,(Min).....	0.4	%
Salt ,(Min).....	0.4	%
Salt ,(Max).....	0.5	%
Sodium ,(Min).....	0.3	%
Sodium ,(Max).....	0.4	%
Vitamin A ,(Min).....	3,623.5	IU/lb
Vitamin D ,(Min).....	53.5	IU/lb
Vitamin E ,(Min).....	11	IU/lb

* Contains not more than 0.9 % protein from non protein nitrogen.

List of Ingredients

Grain products, Processed Grain By-Products, calcium carbonate, plant protein products, salt, urea, potassium chloride, animal fat, magnesium oxide, feed conditioner, wheat red dog, bentonite, monocalcium phosphate, thiamine 40g, vitamin E supplement, vitamin A supplement, soybean oil, vitamin D3 supplement, manganese sulfate, zinc sulfate, copper sulfate, manganous oxide, zinc oxide, sodium selenite, cobalt carbonate, calcium iodate.

Feeding Directions

Feed at a rate of 2 pounds per 100 pounds of body weight per head per day.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: For increased rate of weight gain, feed at a rate of 50 to 200 milligrams Monensin per head per day in not less than 1 pound of feed or, after the 5th day, feed at a rate of 400 milligrams per head per day every other day in not less than 2 pounds of feed. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

Manufactured by:

Lime Ridge Ag Supply, Inc
 115 Minor St
 PO Box 115
 Lime Ridge, WI 53942
 608-986-2626

Texturized
 Net weight 50 lbs (22.67 kg)

MFG DATE: 04/08/2025

MEDICATED