



Lime Ridge Ag Supply Inc
LR RUM HEIF MIN W/FLY#0551

MEDICATED

For beef cattle and/or replacements.

For lactating and non-lactating beef and dairy cows for increased milk production efficiency (production of marketable solid-corrected milk per unit of feed intake). Prevents the development of house flies, stable flies, face flies, and horn flies in the manure of treated cattle.

ACTIVE DRUG INGREDIENT

Monensin (as Monensin Sodium) 1633 g/ton

ACTIVE INGREDIENT

Diflubenzuron 121.6 mg/lb

Guaranteed Analysis

| | | |
|---------------------------|-----------|-------|
| Crude Protein ,(Min)..... | 0.4 | % |
| Crude Fat ,(Min)..... | 0.9 | % |
| Crude Fiber ,(Max)..... | 1.0 | % |
| Calcium ,(Min)..... | 20.5 | % |
| Calcium ,(Max)..... | 21.0 | % |
| Phosphorus ,(Min)..... | 4.4 | % |
| Potassium ,(Min)..... | 0.9 | % |
| Salt ,(Min)..... | 8.0 | % |
| Salt ,(Max)..... | 10.0 | % |
| Sodium ,(Min)..... | 4 | % |
| Sodium ,(Max)..... | 5 | % |
| Vitamin A ,(Min)..... | 198,788.5 | IU/lb |
| Vitamin D ,(Min)..... | 31,185.0 | IU/lb |
| Vitamin E ,(Min)..... | 763 | IU/lb |

List of Ingredients

calcium carbonate, Mono Calcium Phos fr Phos Acid, salt, diflubenzuron, magnesium oxide, dextrose, Vitamin A 30 000 KIU/Kg, soybean oil, Rumensin 90, corn, ethyl acetate, ethyl butyrate, amyl acetate, soy oil, vitamin E supplement, di-alpha tocopheryl acetate, oil, cane molasses, vegetable oil refinery lipid (feed grade), lecithin, phosphoric acid, sulfuric acid, natural and artificial flavoring, propionic acid (preservative) and sodium benzoate (preservative), potassium sulfate, magnesium sulfate, bentonite, sodium selenite, molasses products, vitamin A supplement, vitamin D3 supplement, zinc sulfate, manganous sulfate, copper sulfate, cobalt carbonate, calcium iodate, manganous oxide, zinc oxide, mineral oil, potassium, magnesium, sulfur.

Feeding Directions

Feed at a rate of 4 to 6 ounces per head per day.

CAUTION: Keep out of reach of children. Discard empty container according to local regulations. Never reuse empty container.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions.

Do not feed to lactating goats. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing. A withdrawal time has not been established for pre ruminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: Feed continuously to dry and lactating dairy cows in a total mixed ration. You May Notice (1) Reduced voluntary feed intake in cows fed Monensin. This reduction increases with higher doses of monensin fed. Rule out Monensin as the cause of reduced feed intake before attributing to other causes such as illness, feed management, or the environment. (2) Reduced milk fat percentage in dairy cows fed monensin. This reduction increases with higher doses of Monensin fed. (3) Increased incidence and treatment of cystic ovaries and metritis in dairy cows fed monensin. (4) Reduced conception rates, increased services per animal, and extended days open and corresponding calving intervals in dairy cows fed monensin. Have a comprehensive and ongoing nutritional, reproductive and herd health program in place when feeding Monensin to dairy cows.

Manufactured by:
Lime Ridge Ag Supply, Inc
115 Minor St
PO Box 115
Lime Ridge, WI 53942
608-986-2626

Meal
 Net weight 50 lbs (22.67 kg)

MFG DATE: 07/22/2025

MEDICATED



Lime Ridge Ag Supply Inc
LR RUM HEIF MIN#0503

MEDICATED

For milking dairy cattle and/or replacements.
 Cattle (pastured slaughter, stocker and feeder cattle, and dairy and beef replacement heifers). For increased rate of weight gain .

ACTIVE DRUG INGREDIENT
 Monensin 1674 g/ton

Guaranteed Analysis

| | | |
|---------------------------|-----------|-------|
| Crude Protein ,(Min)..... | 0.4 | % |
| Crude Fat ,(Min)..... | 1.0 | % |
| Crude Fiber ,(Max)..... | 1.0 | % |
| Calcium ,(Min)..... | 21.6 | % |
| Calcium ,(Max)..... | 22.2 | % |
| Phosphorus ,(Min)..... | 4.5 | % |
| Potassium ,(Min)..... | 0.9 | % |
| Salt ,(Min)..... | 8.2 | % |
| Salt ,(Max)..... | 10.2 | % |
| Sodium ,(Min)..... | 4 | % |
| Sodium ,(Max)..... | 5 | % |
| Vitamin A ,(Min)..... | 203,885.5 | IU/lb |
| Vitamin D ,(Min)..... | 31,984.5 | IU/lb |
| Vitamin E ,(Min)..... | 782 | IU/lb |

List of Ingredients

calcium carbonate, Mono Calcium Phos fr Phos Acid, salt, magnesium oxide, dextrose, Vitamin A 30 000 KIU/Kg, soybean oil, Rumensin 90, corn, ethyl acetate, ethyl outyrate, amyl acetate, soy oil, vitamin E supplement, di-alpha tocopheryl acetate, oil, cane molasses, vegetable oil refinery lipid (feed grade), lecithin, phosphoric acid, sulfuric acid, natural and artificial flavoring, propionic acid (preservative) and sodium benzoate (preservative), potassium sulfate, magnesium sulfate, bentonite, sodium selenite, molasses products, vitamin A supplement, vitamin D3 supplement, zinc sulfate, manganous sulfate, copper sulfate, cobalt carbonate, calcium iodate, manganous oxide, zinc oxide, mineral oil, potassium, magnesium, sulfur.

Feeding Directions

Feed at a rate of 4 to 6 ounces per head per day.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal time has not been established for preruminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: For increased rate of weight gain, feed at a rate of 50 to 200 milligrams Monensin per head per day in not less than 1 pound of feed or, after the 5th day, feed at a rate of 400 milligrams per head per day every other day in not less than 2 pounds of feed. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

Manufactured by:
Lime Ridge Ag Supply, Inc
 115 Minor St
 PO Box 115
 Lime Ridge, WI 53942
 608-986-2626

Meal
 Net weight 50 lbs (22.67 kg)

MFG DATE: 07/22/2025

MEDICATED