



Lime Ridge Ag Supply Inc
LR STEER MIN R2000 #0517

MEDICATED

Mature reproducing beef cows; For improved feed efficiency; for prevention and control of coccidiosis due to Eimeria bovis and E. zuernii.

ACTIVE DRUG INGREDIENT
 Monensin Sodium 1995 g/ton

Guaranteed Analysis

Crude Protein ,(Min).....	2.0	%
Crude Fat ,(Min).....	2.3	%
Crude Fiber ,(Max).....	1.0	%
Calcium ,(Min).....	23.4	%
Calcium ,(Max).....	24.0	%
Phosphorus ,(Min).....	2.8	%
Salt ,(Min).....	11.7	%
Salt ,(Max).....	13.7	%
Sodium ,(Min).....	6	%
Sodium ,(Max).....	7	%
Vitamin A ,(Min).....	147,693.0	IU/lb
Vitamin D ,(Min).....	12,491.5	IU/lb
Vitamin E ,(Min).....	102	IU/lb

List of Ingredients

calcium carbonate, salt, Mono Calcium Phos fr Phos Acid, Processed Grain By-Products, dextrose, soybean oil, Rumensin 90, roughage products, vitamin A supplement, mineral oil, potassium sulfate, magnesium sulfate, bentonite, vitamin E supplement, sodium selenite, molasses products, vitamin D3 supplement, zinc sulfate, manganous sulfate, copper sulfate, cobalt carbonate, calcium iodate, manganous oxide, zinc oxide.

Feeding Directions

Feed at a rate of 4 to 6 ounces per head per day.

CAUTION: Do not allow horses or other equines access to formulation containing Monensin. Ingestion of Monensin by equines has been fatal. Monensin medicated cattle and goat feed is safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Feeding undiluted or mixing errors resulting in high concentrations of Monensin has been fatal to cattle and could be fatal to goats. Must be thoroughly mixed in feeds before use. Do not feed undiluted. Do not exceed the levels of Monensin recommended in the feeding directions as reduced average daily gains may result. If feed refusals containing Monensin are fed to other groups of cattle, the concentration of Monensin in the refusals and amount of refusals fed should be taken into consideration to prevent Monensin overdosing.

WARNING: A withdrawal time has not been established for prurminating calves. Do not use in calves to be processed for veal.

LIMITATIONS: For increased rate of weight gain, feed at a rate of 50 to 200 milligrams Monensin per head per day in not less than 1 pound of feed or, after the 5th day, feed at a rate of 400 milligrams per head per day every other day in not less than 2 pounds of feed. For prevention and control of coccidiosis, feed at a rate of 0.14 to 0.42 milligram per pound of body weight per day, depending on severity of challenge, up to 200 milligrams per head per day. During first 5 days of feeding, cattle should receive no more than 100 milligrams per day in not less than 1 pound of feed.

Manufactured by:
 Lime Ridge Ag Supply, Inc
 115 Minor St
 PO Box 115
 Lime Ridge, WI 53942
 608-986-2626

Meal
 Net weight 50 lbs (22.67 kg)

MFG DATE: 07/22/2025

MEDICATED